

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>A61K 7/00</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 98/49999</b>
		<b>(43) International Publication Date:</b> 12 November 1998 (12.11.98)
<b>(21) International Application Number:</b> PCT/EP98/02795 <b>(22) International Filing Date:</b> 4 May 1998 (04.05.98)  <b>(30) Priority Data:</b> 97201304.9      2 May 1997 (02.05.97)      EP <b>(34) Countries for which the regional or international application was filed:</b> AT et al.  <b>(71) Applicant (for all designated States except US):</b> GIST-BROCADES B.V. [NL/NL]; Wateringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> LAMBERS, Johannes, Wilhelmus, Jacobus [NL/NL]; Altena 10, NL-2641 LB Pijnacker (NL). STREEKSTRA, Hugo [NL/NL]; Water- ingstraat 28 I, NL-1017 SP Amsterdam (NL).  <b>(74) Agents:</b> VISSER-LUIRINK, Gesina et al.; Gist-Brocades B.V., Patents and Trademarks Dept., Wateringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL).		<b>(81) Designated States:</b> BR, JP, KR, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> ANTIMICROBIAL COMPOSITIONS FOR TOPICAL USE		
<b>(57) Abstract</b>		
<p>The present invention discloses that topically occurring microbial growth is inhibited by applying a topical composition comprising a sphingoid base. Specifically, said sphingoid base is effectively formulated in combination with a surfactant.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NI	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## ANTIMICROBIAL COMPOSITIONS FOR TOPICAL USE

### Field of the invention

5           The present invention relates to the field of topical compositions comprising sphingoid bases functioning as antimicrobial agents.

### Background of the invention

10           Healthy human skin is colonized by a number of different microorganisms responsible for maintaining the natural microbial equilibrium of the skin. However, outgrowth of certain species within the skin microflora can easily occur, thereby causing cosmetically and dermatologically undesirable phenomena, like unpleasant body odour, and in a worse case,  
15           infections. Several skin conditions are known which are associated with unwanted microbial growth. For instance, wounded or diseased skin especially is prone to superinfections by *Staphylococcus aureus*, a bacterium which also is the most important infective agent in patients suffering from atopic eczema. In addition, acne is associated with outgrowth of the  
20           bacterium *Propionibacterium acnes*. Also, fungal skin infections (mycoses) are known to occur frequently.

          It has been known for several decades that skin surface lipids contain one or more lipid compounds possessing antimicrobial activity against gram-positive bacteria (Burtenshaw (1942), J. Hyg. 42, 184-209). These  
25           antimicrobial lipid compounds were thought to be mainly the free fatty acids released from sebaceous triglycerides by lipases from the normal microflora (Kearney *et al.* (1984), Br. J. Dermatol. 110, 593-599). Only recently, attention has been focused on the role of sphingolipids in this respect.

          In a recent series of papers, Bibel *et al.* attribute an antimicrobial  
30           activity to sphingoid bases. In a first paper, test mixtures containing about 0.0005 to 0.005 % of a sphingoid base were described to inhibit *in vitro* microbial growth (Bibel *et al.* (1992), J. Invest. Dermatol. 98, 269-273).

However, high concentrations of ethanol were additionally present in these mixtures and ethanol is known as a solvent which itself substantially contributes to antimicrobial activity. Furthermore, no or a very low activity against gram-negative bacteria was observed. In *in vivo* experiments (Bibel *et al.* (1992), *supra*; Bibel *et al.* (1995), Clin. Exper. Dermatol. 20, 395-400), sphingoid bases were applied in much higher concentrations, i.e. as an 1.6% ethanolic solution or as a suspension of an 1.5% ethanolic solution in petrolatum. However, these formulations are not considered to enable an effective delivery of an active ingredient to the skin. For instance, Bibel *et al.* (1995) reported granulation of sphinganine upon drying, resulting in a decreased availability of this compound.

The present invention discloses compositions for the inhibition of topically-occurring microbial growth which comprise an effectively formulated sphingoid base, i.e. a sphingoid base which is formulated without ethanol in a high concentration.

### Description of the invention

The present invention discloses that sphingoid bases have a potent antimicrobial activity in the absence of inhibiting concentrations of an antimicrobial solvent like ethanol. Effective formulations are disclosed which are suitable for topical application on various skin conditions associated with undesired microbial growth.

The present invention further discloses that the concentration of a sphingoid base necessary to obtain a substantial antimicrobial effect *in vitro* in the absence of inhibiting concentrations of ethanol should be at least about 0.005 wt %. When higher concentrations are used, i.e. 0.01, 0.02, 0.04 or 0.08 wt %, the antimicrobial effect of a sphingoid base increases.

Throughout the invention, the terms "antimicrobial activity / effect" or "growth-inhibitory activity / effect" are used synonymously.

Microorganisms which are susceptible to the antimicrobial activity of sphingoid bases include bacteria, yeasts and fungi.

When using a concentration of at least about 0.005 % of a sphingoid base, the present invention discloses that sphingoid bases also display a growth-inhibitory activity on gram-negative bacteria. In particular, the present invention discloses that sphingoid bases display a growth-inhibitory activity against gram-negative bacteria which is about similar as compared to the activity against gram-positive bacteria.

The present invention further discloses that sphingoid bases display a substantial antifungal activity. An antifungal activity is understood to include a growth-inhibitory activity on yeasts as well as on filamentous fungi.

The present invention discloses that sphingoid bases included in compositions for topical use are effectively formulated when combined with a surfactant selected from the group of ionic (anionic and/or cationic) and nonionic surfactants. Preferably, the surfactant is selected from the group of nonionic surfactants, more preferably from the group of ethoxylated-sorbitan-esters, such as Tween 80.

The topical compositions according to the invention include compositions wherein water is used as a solvent, compositions wherein an emollient (e.g. a fat or an oil) is used as a solvent and compositions where water and a fat or an oil are used (emulsions) as a solvent.

The concentration in which the sphingoid base is effectively formulated in a topical composition may range from 0.001 to 5 wt %, preferably from 0.005 to 5 wt %, more preferably from 0.01 to 2.5 wt %, most preferably from 0.02 to 1 wt %, especially preferably from 0.02 to 0.5 wt %.

It thereby may depend on the type of application which concentration of a sphingoid base advantageously is used. Typically, treatment of an infection, e.g. a wound infection, may require a higher dose of a sphingoid base than preventive use, e.g. the normalisation of skin flora.

The surfactant typically is applied in a concentration ranging from 0.01 to 10 %, preferably from 0.1 to 5 %, more preferably from 0.5 to 2.5 %.

The type of sphingoid base used is not critical to the invention. Typically, a sphingoid base is selected from the group of sphingamines,

sphingosines or phytosphingosines. Preferably, a sphingoid base is selected from the group of phytosphingosines.

The sphingoid bases used according to the invention may be obtained from any suitable source, e.g. from a natural source or from a chemical synthesis process. However, it is desirable to apply a production process such that a sphingoid base is obtainable in sufficient quantities at commercially feasible prices. In that regard, some current sources of sphingoid bases may have disadvantages. In case of chemical synthesis, it is very difficult to prepare the correct stereochemical configuration. In case of purification of animal and/or plant tissue extracts, the amounts of sphingoid bases are very small, making their isolation costly. Moreover, animal sources are believed to be unsafe due to the presence of viruses and other infectious agents, such as the agent causing BSE (mad cow's disease).

Therefore, sphingoid bases are preferably obtained from a microbial fermentation process. More preferably, they are obtained from a yeast, especially preferably from *Pichia ciferrii*. In one embodiment of the invention, the sphingoid base phytosphingosine is obtained from *Pichia ciferrii*-derived tetraacetyl-phytosphingosine (TAPS), by a suitable deacetylation reaction. The deacetylation may be chemical, e.g. by base catalyzed hydrolysis with potassium hydroxide, or enzymatical. After alkaline hydrolysis of TAPS, the resulting phytosphingosine may be purified. Such a purification can occur by any method known to a person skilled in the art. Yeast-derived phytosphingosine is human skin-identical, as it is reported to have the same stereochemical configuration as mammalian phytosphingosine, i.e. the D-D-erythro configuration.

Compositions according to the invention comprising a sphingoid base are suitable for topical application, whereby topical application is understood to comprise cosmetic and/or dermatological application on the skin, on hair and on the epithelial linings of mouth, nose, eye, urogenital tract, and the like. Topical compositions including a sphingoid base are suitable to apply for various topically occurring undesirable and/or abnormal conditions associated with microbial activity.

Topical compositions according to the invention comprising a sphingoid base also advantageously are applied in the form of a plaster, dressing, and the like.

Examples of topically occurring undesirable and/or abnormal conditions in which topical compositions comprising a sphingoid base are advantageously applied are: acne, dandruff, mouth and/or lip infections, mycoses, various other skin-infectious diseases or vaginal infections. Topical compositions comprising a sphingoid base are further advantageously applied for wound-healing, e.g. in case of burns, and for normalisation of skin flora.

Due to their antimicrobial activity, sphingoid bases additionally may function as a preservative in cosmetic and dermatological compositions, to decrease and/or substitute for existing chemical preservatives.

In the following examples, the antibacterial and antifungal activity of sphingoid bases is shown. In addition, various examples of effective formulations suitable for topical application of a sphingoid base are given.

#### Description of the figures

Figure 1. CFUs obtained after incubation of *Staphylococcus aureus* and *Corynebacterium xerosis* with increasing concentrations of sphingosine.

Figure 2. CFUs obtained after incubation of *Escherichia coli* during an increasing time period with three concentrations of phytosphingosine.

Figure 3. CFUs after incubation of *Pseudomonas aeruginosa* during an increasing time period with two concentrations of phytosphingosine.

Figure 4. CFUs after incubation of *P. acnes* during an increasing time period with three concentrations of phytosphingosine.

**Experimental****Strains**

Gram-positive bacteria:

5	<i>Micrococcus luteus</i>	ATCC 9341	normal flora
	<i>Staphylococcus aureus</i>	ATCC 9196	normal flora, boils
	<i>Corynebacterium xerosis</i>	A 2150	sweat
	<i>Propionibacterium acnes</i>	ATCC 6919	acne

10 Gram-negative bacteria:

	<i>Pseudomonas aeruginosa</i>	ATCC 9027	wound infections
	<i>Escherichia coli</i>	ATCC 11229	

Yeasts and fungi:

15	<i>Saccharomyces cerevisiae</i>	ATCC 9763	
	<i>Candida albicans</i>	ATCC 10231	
	<i>Microsporum canis</i>	CBS 283.63	

**Measurement of antibacterial activity**

20 All bacterial strains were grown overnight in a 500 ml shake flask containing 100 ml Brain Heart Infusion medium, at 37 °C and 280 rpm. Only *P. acnes* was grown anaerobically in BHI medium flushed with sterile nitrogen gas, for 48-72 hours at 30 °C without agitation.

25 The antibacterial activity of a sphingoid base of choice is measured as the amount of colony forming units (CFUs) obtained after incubation of a bacterial strain with a sphingoid base. To this end, 100  $\mu$ l of a 10 times diluted overnight culture in 1% Neopeptone were added to 300  $\mu$ l of a solution containing 0.7 % Neopeptone, 0.6 % ethanol, 1.1 % Tween 80 and an appropriate amount of a sphingoid base. The resulting mixture was  
30 incubated at 37 °C during a specified time period, whereupon 100  $\mu$ l of a 10<sup>6</sup> times dilution in physiological salt were plated onto BHI agar plates. The resulting colonies were counted.



### Measurement of antifungal activity

#### Liquid dilution method in 96-wells microtiterplates

- a stock solution of 10 mg of phytosphingosine/ml was prepared in water containing 20 % ethanol and 16 % Tween 80;
- 5 - from the stock solution a ten-fold dilution was prepared in Sabouraud dextrose broth (Difco);
- from *Saccharomyces cerevisiae* and *Candida albicans* cultured on Sabouraud dextrose broth several serial dilutions were prepared in said medium in a total count twice as much as is needed in the test;
- 10 - 100  $\mu$ l of the ten-fold dilution of phytosphingosine was added to the first wells of the left side (in a vertical way) of a 96-wells microtiterplate;
- to each row of wells 100  $\mu$ l of the appropriate culture and dilution was added in such a way that the highest total count is on top;
- after mixing the first well, 100  $\mu$ l from this well was added to the second
- 15 - well at the right side;
- after mixing the second well, 100  $\mu$ l from this well was added to the third well at the right side etc.;
- for each row this dilution rate was made, thus preparing from left to right a dilution rate of 1000, 500, 250 microgram of phytosphingosine/ml etc.
- 20 - and from top to bottom a serial dilution rate in total count;
- after incubation at appropriate temperature and time, growth was measured by examining sedimentation on the bottom of the wells using a mirror;
- the concentration of phytosphingosine in the first well where no growth
- 25 - was observed, is the value for the minimal inhibitory concentration (mic).

#### Agar dilution method

- a stock solution of 20 mg of phytosphingosine/ml was prepared in water containing 20 % ethanol and 16 % Tween 80;
- 30 - from the stock solution a twenty-fold dilution was prepared in molten and cooled down Malt Extract Agar (MEA, Difco);
- 50 ml of this dilution was mixed with 50 ml of molten and cooled down MEA;

- 50 ml of the second dilution was mixed with 50 ml of molten and cooled down MEA etc, creating a dilution rate of 1000, 500, 250  $\mu$ g phytosphingosine/ml MEA etc.;
- from each dilution 20 ml was poured into plastic petridishes (9 cm) and cooled down to harden;
- *Microsporum canis* was cultured on MEA at appropriate temperature and time;
- after culturing sterile glass beads in physiological salt were added;
- the culture was shaken till a homogeneous mass was obtained;
- 10  $\mu$ l of the thus obtained cultures were spotted onto the various agardishes and cultured at appropriate temperature and time;
- from cultured *Saccharomyces cerevisiae* in Sabouraud dextrose broth, appropriate serial dilutions were made in the same broth;
- 10  $\mu$ l of the thus obtained cultures were spotted onto the various agardishes and cultured at appropriate temperature and time;
- after incubation the growth was visually examined;
- the concentration of phytosphingosine in the first plate where no growth was observed, is the value for the minimal inhibitory concentration (mic).

The sphingoid bases used were sphingosine (Sigma) and phytosphingosine (obtained from deacylation of *Pichia ciferri*-derived tetraacetylphytosphingosine).

### Example 1

#### Antibacterial activity of various concentrations of sphingosine

The antibacterial activity of sphingosine (S) on two bacterial strains was analyzed by incubating a bacterial strain with increasing concentrations of sphingosine for 60 minutes at 37 °C (see Experimental for further details).

Two representative graphs, depicting the antibacterial activity of sphingosine against *S. aureus* and *C. xerosis* (Figure 1), show that the amount of CFUs decreases with an increasing concentration of sphingosin from 0.005 to 0.02%.

A comparable antibacterial activity of sphingosine and/or phytosphingosine was measured using *M. luteus*, *E. coli*, *P. aeruginosa* and *P. acnes*.

### Example 2

#### Antibacterial activity of phytosphingosine as related to time

The antibacterial activity of phytosphingosine (PS) on various bacteria as related to time was analyzed by incubating a selected bacterial strain with a solution containing phytosphingosine at 37 °C during a time period up to 240 minutes (see Experimental for further details).

The results of the incubation of *E. coli*, *P. aeruginosa* and *P. acnes* with two or three different concentration of phytosphingosine are depicted in Figures 2, 3 and 4, respectively. It is shown that the outgrowth of bacteria which occurred at longer incubation times is prevented by using higher concentrations of phytosphingosine, i.e. 0.02 to 0.08 wt %.

### Example 3

#### Antifungal activity of phytosphingosine

20

Antifungal activity of phytosphingosine on the yeasts *S. cerevisiae* and *C. albicans* was determined using the liquid dilution method (see Experimental). The m.i.c. values obtained are indicated below. These values indicate that phytosphingosine has a potent growth inhibitory activity on yeasts.

25

#### *Saccharomyces cerevisiae*

	<u>m.i.c</u>
0.7x10 <sup>6</sup> cfu/ml	60 ppm
0.7x10 <sup>5</sup> cfu/ml	15 ppm
0.7x10 <sup>4</sup> cfu/ml	15-30 ppm
0.7x10 <sup>3</sup> cfu/ml	4-8 ppm
0.7x10 <sup>2</sup> cfu/ml	2-8 ppm

30

35

*Candida albicans*

		<u>m.i.c</u>
	0.9x10 <sup>6</sup> cfu/ml	125-500 ppm
	0.9x10 <sup>5</sup> cfu/ml	60-125 ppm
5	0.9x10 <sup>4</sup> cfu/ml	125-250 ppm
	0.9x10 <sup>3</sup> cfu/ml	15-60 ppm
	0.9x10 <sup>2</sup> cfu/ml	125 ppm

Antifungal activity of phytosphingosine was also determined using a  
 10 dermatophyt. (*M. canis*). This was done using the agar dilution method (see  
 Experimental). For comparison, *S. cerevisiae* was also included. The results  
 are indicated in the table below. It is clear that phytosphingosine also is able  
 to significantly inhibit the growth of a fungus considered to be a  
 representative of a skin fungus.

15

concentration ppm	<i>Sacch. cerevisiae</i> 10 <sup>3</sup> cfu/ml	<i>Microsporum canis</i>
1,000	- / -	- / -
500	- / -	pp / -
20 250	- / -	- / -
125	- / -	- / -
60	- / -	- / pp
30	- / -	+ / +
15	+ + + / + + +	+ / +
25 7.5	+ + + / + + +	+ / +
0	+ + + / + + +	+ + + / + + +

+ + + = good growth, + + = moderate growth,  
 + = very moderate growth, pp = pin point colony, - = no growth

30

**Example 4****Compositions comprising a non-ionic surfactant****Standardized solution**

5	Non-ionic surfactant	1-15%
	Phytosphyngosine	0.1-2%
	Humectant	5-10%
	Water	qs 100

**Anti acne skin cleansing lotion**

10	<b>A.</b>	PPG-26-buteth- 26 +	
		PEG40 hydrogenated castor oil	1%
		Phytosphyngosine	0.2%
		Butylene glycol	5%
15	<b>B.</b>	Water	qs 100

**Procedure:** Combine all ingredients of A by mixing. Add B to A while stirring and keep on stirring until the product is homogeneous.

**Antimicrobial mouthwash concentrate**

20	<b>A.</b>	Glycerin	20%
		Phytosphyngosine	0.1%
		Polysorbate 80	1%
25	<b>B.</b>	Flavor	1.5%
		Sodium Saccharin	0.03%
		Water	qs 100

**Procedure:** Dissolve phytosphyngosine into the polysorbate at room  
30 temperature, add the glycerin. Then add B to A and mix until homogeneous.

**Example 5**Oil/water emulsions**Standardized emulsion**

5	A.	various oil phase	10-25%
		non-ionic surfactants	2-7%
	B.	humectant	5-10%
		non-ionic surfactant	1-4%
10		phytosphingosine	0.1-1%
		water	qs100

**Therapeutic Foot cream**

	A.	Water	70%
15	B.	Mineral oil	5%
		C12-C15 alcohol benzoate	2%
		Propylene glycol dicapralate	5%
		Dimethicone	1%
		Ceteareth 21	3%
20		Ceteareth 2	2%
	C.	Propylene glycol	10%
		Ethoxylated hydrogenated castor oil	1%
		Phytosphingosine	0.2%
25	D.	Perfume	qs

Procedure : Heat A and B to 75-80°C and mix for 30 minutes with good agitation. Cool to 40°C add C and D.

30

**Example 6**Cleansing compositions**Standardized soap solution**

35	Anionic surfactant	0-15%
	Amphoteric surfactant	0-15%
	Non-ionic surfactant	0-20%
	Phytosphingosine	0.1-1%
	EDTA	0.05%
40	Water	qs100

**Anti-microbial facial cleanser**

5	<b>A.</b>	Sodium laureth sulfate	12%
		Cocamidopropylbetaine	3%
		Cocamide DEA	2%
		Phytosphingosine	0.1%
	<b>B.</b>	Propylene glycol	20%
		Water	qs 100
10	<b>C.</b>	Fragrance	qs

**Procedure:** Combine A and B at 40°C. When homogeneous add C under stirring.

**15 Skin Lotion with phytosphingosine**

15	<b>A.</b>	Water	70%
		Carbomer	0.3%
20	<b>B.</b>	Mineral oil	5%
		C12-C15 alcool benzoate	2%
		Propylene glycol dicapralate	5%
		Cetyl alcohol	1%
		Stearic acid	3%
25	<b>C.</b>	Dimethicone	1%
	<b>D.</b>	Butylene glycol	7%
		Oleic acid	1%
		Phytosphingosine	0.2%
30	<b>E.</b>	Triethanolamine	1%
		Water	qs100
	<b>F.</b>	Perfume	qs

35

**Procedure:** Disperse A. Heat B to 75-80°C and mix until uniform. Combine with A and mix for 30 minutes with good agitation. Cool to 60°C and add C. Cool to 40°C add D , E and F.

40

**Example 7****Gel composition****Standardized gel formulation**

5	Gellifiant	0.5-3%
	Humectant	5-50%
	Phytosphingosine	0.1-2%
	Water	qs100

**10 Anti acne gel**

A.	Natrosol	2,25%
	Water	qs100

B.	Ethoxydiglycol	20%
15	Phytosphingosine	0.2%

**Procedure:** Disperse the Carbomer into the water, then add B. Adjust the pH with triethylamine (TEA).

**20 Roll on deodorant**

A.	Propylene glycol	30%
	Phytosphingosine	0.1%

B.	Natrosol	1.25%
25	EDTA	0.025%
	Water	qs 100

**Procedure:** Disperse the Natrosol into the water, then add A.



Gist-brocades B.V.

## CLAIMS

- 5           1. Use of a sphingoid base for the manufacture of a topical composition comprising a surfactant selected from the group of nonionic and ionic surfactants for the inhibition of topically-occurring microbial growth.
2. The use of claim 1, wherein the topical composition comprises a  
10   surfactant selected from the group of nonionic surfactants.
3. The use of claim 1 or 2, wherein said microbial growth is caused by a microorganism selected from the group of bacteria, yeasts and fungi.
- 15          4. The use of claim 3, wherein the bacteria are gram-negative bacteria.
5. The use of claim 1 to 4, wherein the topical composition comprises a sphingoid base in a concentration ranging from 0.001 to 5 wt%,  
20   preferably from 0.005 to 5 wt%, more preferably from 0.01 to 2.5 wt%, most preferably from 0.02 to 1 wt%, especially preferably from 0.02 to 0.5 wt%.
6. The use of claim 1 to 5, wherein the sphingoid base is selected  
25   from the group of sphinganine, sphingosine and phytosphingosine.
7. The use of claim 6, wherein the sphingoid base is phytosphingosine.
- 30          8. A method for inhibiting topically-occurring microbial growth, comprising application of a topical composition comprising a sphingoid base and a surfactant selected from the group of nonionic and ionic surfactants.

9. The method of claim 8, wherein the topical composition comprises a sphingoid base in a concentration ranging from 0.001 to 5 wt%, preferably from 0.005 to 5 wt%, more preferably from 0.01 to 2.5 wt%, most preferably from 0.02 to 1 wt%, especially preferably from 0.02 to 0.5 wt%.

5

10. The method of claim 8 or 9, wherein the sphingoid base is selected from the group of sphingamines, sphingosines and phytosphingosines.

10 11. The method of claims 8 to 10, wherein the sphingoid base is phytosphingosine.

12. The method of claims 8 to 11, wherein the topical composition is a cosmetic composition.

15




Figure 1




Figure 2




Figure 3

4/4








Figure 4



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 7/48, 7/32, 7/22, A01N 33/08</b>		<b>A3</b>	(11) International Publication Number: <b>WO 98/49999</b>
(21) International Application Number: PCT/EP98/02795		(43) International Publication Date: 12 November 1998 (12.11.98)	
(22) International Filing Date: 4 May 1998 (04.05.98)		(81) Designated States: BR, JP, KR, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(30) Priority Data: 97201304.9 2 May 1997 (02.05.97) EP		Published With international search report.	
(34) Countries for which the regional or international application was filed: AT et al.		(88) Date of publication of the international search report: 4 February 1999 (04.02.99)	
(71) Applicant (for all designated States except US): GIST-BROCADES B.V. [NL/NL]; Wateringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL).			
(72) Inventors; and (75) Inventors/Applicants (for US only): LAMBERS, Johannes, Wilhelmus, Jacobus [NL/NL]; Altena 10, NL-2641 LB Pijnacker (NL). STREEKSTRA, Hugo [NL/NL]; Water- ingstraat 28 I, NL-1017 SP Amsterdam (NL).			
(74) Agents: VISSER-LUIRINK, Gesina et al.; Gist-Brocades B.V., Patents and Trademarks Dept., Wateringseweg 1, P.O. Box 1, NL-2600 MA Delft (NL).			

## (54) Title: ANTIMICROBIAL COMPOSITIONS FOR TOPICAL USE



## (57) Abstract

The present invention discloses that topically occurring microbial growth is inhibited by applying a topical composition comprising a sphingoid base. Specifically, said sphingoid base is effectively formulated in combination with a surfactant.

**ATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NI	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakistan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 98/02795

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 6 A61K7/48 A61K7/32 A61K7/22 A01N33/08

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 A61K A01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DE 195 03 423 A (BEIERSDORF) 8 August 1996 see examples 5, 14-18 ---	1-12
X,P	DE 196 43 585 A (BEIERSDORF) 23 April 1998 see page 4, line 42; claims 1-6 ---	1-12
A	STN, File Supplier, Karlsruhe, DE, File XP002083142 cited in the application Medline, AN=96169713 see the abstract ---	1-12
A	STN, File Supplier, Karlsruhe, DE, File XP002083143 Medline, AN=94152179 see the abstract --- -/--	1-12

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"S" document member of the same patent family

Date of the actual completion of the international search

4 November 1998

Date of mailing of the international search report

17/11/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Fischer, J.P.

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/02795

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	STN, File Supplier, Karlsruhe, DE, File XP002083144 Medline, AN=93099560 see the abstract ---	1-12
A	STN, File Supplier, Karlsruhe, DE, File XP002083145 cited in the application Medline, AN=92185246 see the abstract -----	1-12

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/EP 98/02795

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19503423 A	08-08-1996	WO 9623479 A EP 0806935 A	08-08-1996 19-11-1997
DE 19643585 A	23-04-1998	NONE	

**THIS PAGE BLANK (USPTO)**

DOCKET NO: RWS-80019

SERIAL NO: 09/936, 316

APPLICANT: Streekstra et al.

LERNER AND GREENBERG P.A.

P.O. BOX 2480

HOLLYWOOD, FLORIDA 33022

TEL. (954) 925-1100